

THE UNITED SHAYES OF AMERICA

TO ALL TO WHOM THESE: PRESENTS: SHALL COME:

Syngenta Seeds, Inc.

MICCERS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY, PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE IGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR PROPAGATION, OR STOCKING IT FOR ANY OF THE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN, FIELD

'NP2467'

In Vestimonn Marrest, I have hereunto set my hand and caused the seal of the Mant Anxiety Protection Office to be affixed at the City of Washington, D.C. this twenty-third day of November, in the year two thousand and seven.

Aus

Benze

Commissioner Plant Variety Protection Offic Agricultural Marketing Servi of Agriculture

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE

Application is required in order to determine if a plant

(Instructions and information collection burden s	tatement on rever	se)	T				
			2. TEMPORARY DESIGNATION EXPERIMENTAL NAME	ON OR 3	3. VARIETY NAME		
Syngenta Seeds, Inc.					NP2467		
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code	e, and Country)		5. TELEPHONE (include area		FOR OFFICIAL USE ONLY		
P.O. Box 959			(763) 593-733	33	VPO NUMBER		
Minneapolis, MN 55440			6. FAX (include area code)	7	00500100		
			(763) 593-78	28	ILING DATE		
7. IF THE OWNER NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZATION (corporation, partnership, association, etc.)	8. IF INC	CORPORATED, GIVE E OF INCORPORATION	9. DATE OF INCORPORATIO	N	Jan. 21,2005		
Corporation		laware	Sept. 24, 197	6	·		
10 NAME AND ADDRESS OF OWNER REPRESENTATIVE(S) TO SE	RVE IN THIS APPLICA	TION. (First person listed will re	ceive all papers)		FILING AND EXAMINATION FEES:		
	<u>.</u> .	h /		Ę			
Đavid L. Young	S. Rewold	dt RO Roy	(500	E	DATE 1/21/05		
Syngenta Seeds, Inc. 2369	33044 57	dt reet, P.O. Box 244		R			
317-330th-Street Slater	+ 4 30.	244		E V	s 768.00		
-Stanton, MN 55018-4308				Š	1		
	-			-	DATE 10 1 07		
11. TELEPHONE (Include area code) 5/5 - 685 - 520 / 5/5 - 685 - 520 / 5/5 - 685 - 520 / 5/5 - 68		_	t@syngenta.com		IND (Common Name)		
(507) 663-7620 (507) 645	5-7519	-dave.young@	syngenta.com	Col	rn (dent)		
15 GENUS AND SPECIES NAME OF CROP		16. FAMILY NAME (Bolanic	1	17. IS THE THE PRINT	VARIETY A FIRST GENERATION		
Zea mays L.		Graminea	e		YES NO		
 CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITT reverse) 	TED (Follow instructions	on 19. DOES THE C	OWNER SPECIFY THAT SEED OF SEED? See Section 83(a) of the section 100 (a) (b) (c) (c) (c) (c) (c) (d) (d) (d) (d) (d) (d) (d) (d) (d) (d	F THIS VARIE	TY BE SOLD AS A CLASS OF		
a. X Exhibit A. Origin and Breeding History of the Variety b. X Exhibit B. Statement of Distinctness		,	fES (if "yes", answeritems 20 and 21 below)	X	NO (If "no," go to item 22)		
b.		20. DOES THE CO	OWNER SPECIFY THAT SEED O	F THIS VARIE	TY BE LIMITED AS TO NUMBER		
 d. Exhibit D. Additional Description of the Variety (Optional) e. Exhibit E. Statement of the Basis of the Owner's Owners! 		I ' ' ' <u> </u>	ES		NO		
Voucher Sample (2,500 viable untreated seeds or, for tub verification that lissue culture will be depositied and maint	er propagated varieties		ITEM 20, WHICH CLASSES OF I	RODUCTION	BEYOND BREEDER SEED?		
repository) g. Filing and Examination Fee (\$2,450), made payable to "Ti		I	FOUNDATION REGISTERED CERTIFIED				
States" (Mail to the Plant Variety Protection Office) 22. HAS THE VARIETY (INCLUDING ANY HARVESTED MATERIAL) O	D A LIVEDIN SOANISC	ED 22 16 TUE VADI	23. IS THE VARIETY OR ANY COMPONENT OF THE VARIETY PROTECTED BY INTELLECTUAL				
FROM THIS VARIETY BEEN SOLD, DISPOSED OF, TRANSFERRI OTHER COUNTRIES?	ED, OR USED IN THE I	I. S. OR PROPERTY	PROPERTY RIGHT (PLANT BREEDER'S RIGHT OR PATENT)?				
YES NO IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALE, DISPO	OSITION, TRANSFER, (OR USE IF YES, PLEA	IF YES, PLEASE GIVE COUNTRY, DATE OF FILING OR ISSUANCE AND ASSIGNED				
FOR EACH COUNTRY AND THE CIRCUMSTANCES. (Please use 24. The owners declare that a viable sample of basic seed of the variety			NUMBER. (Please use space in				
7-1 This divites a secure trial a value sample to lead to use valuely for a tuber propagated variety a tissue culture will be deposited in a The undersigned owner(s) is(are) the owner of this sexually reprodu-	public repository and m	aintained for the duration of the	certificate.	_			
and is entitled to protection under the provisions of Section 42 of the Owner(s) is(are) informed that false representation herein can jeopa	Plant Variety Protection	A Act.			and the second and an experience to the second		
SIGNATURE OF OWNER	//	SIGNATURE OF	OWNER				
Karald Sterms							
NAME (Please print or type)		NAME (Please pri	int or type)				
Ronald S. Ferriss, Ph.D.							
CAPACITY OR TITLE Director, Strategy Facilitation	DATE	CAPACITY OR TI	TLE		DATE		

INSTRUCTIONS

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$2,450 (\$300 filing fee and \$2,150 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 500, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$300 for issuance of the certificates. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvp.htm

ITEM

18a, Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively:
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 22. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 5.5 of the Act for instructions on claiming the benefit of an earlier filing date.

22. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

Released in a Hybrid, (01/23/04) offered for sale in the U.S. and Canada.

23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

U.S. patent filed on Jan. 20, 2005 (serial # 11/039, 563) issued on Jan. 2, 2007 (patent # 7, 157,631)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the variety names proposed by contacting: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center-East, Beltsville, MD 20705. Telephone: (301) 504-8089.

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Officer, OIRM, AG Box 7630, Jamie L. Whitten Building, Washington, D.C. 20250. When replying, refer to OMB No. 0581-0055 and form number in your letter. Under the PRA of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (traitle, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-2791. To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call (202) 720-7327 (voice) or (202) 720-1127 (TDD). USDA is an equal opportunity employer.

S8T-470 (6-98) designed by the Plant Variety Protection Office with WordPerfect 6.0a. Replaces STD-470 (03-96) which is obsolete.

Syngenta Seeds, Inc. NP2467 Exhibit A

Origin and Breeding History of Corn Inbred Line NP2467

The corn inbred line NP2467 was derived from the initial cross of inbred line NP2158 and inbred line NP2153. Inbred line NP2158 was developed by crossing Holden's Foundation Seeds inbred line LHE136 and inbred line NP2455. NP2455 was derived from a bag of Pioneer 3358 purchased in 1986. NP2153 was derived from B73 and was developed by Illinois Foundation Seeds, Inc.

After the initial cross of NP2158 and NP2153, the breeding method was simple pedigree ear-to-row for the development of inbred line NP2467.

The initial self-pollination of the F1 source was made at Boynton Beach, FL in the Ciba Geigy Company nursery during the winter of 1995. This was followed by self-pollinating and pedigree selection at Bloomington, IL; Boynton Beach, FL; Kaunakakai, HI; and Salinas, PR from 1996 through 2000.

The chronology of inbreeding is given below:

1995 Boynton Beach, FL (Summer)	Initial cross was made to create F ₁ seed.
1995 Boynton Beach, FL (Winter)	F ₁ plants self-pollinated to create F ₂ population.
1996 Kaunakakai, HI (Summer)	F ₂ plants self-pollinated to produce F ₃ families.
1996 Boynton Beach, FL (Winter)	Plants of selected F_3 families self-pollinated to produce F_4 families.
1997 Bloomington, IL	Plants of selected F_4 families self-pollinated to produce F_5 families.
1998 Bloomington, IL	Plants of selected $F_{\rm 5}$ families self-pollinated to produce $F_{\rm 6}$ families.
1999 Bloomington, IL	Plants of selected F_6 families self-pollinated to produce F_7 families.
1999 Salinas, PR (Winter)	Plants of selected F_7 families self-pollinated to produce F_8 lines.
2000 Bloomington, IL	Selected F_8 lines self-pollinated and ear rows bulked to produce Breeder Seed.

Selection criteria used during the inbreeding process included synchronous male and female flowering, plant health, ear fill, pre-harvest intactness and resistance to various stalk rots. Plants within each generation were also closely evaluated for uniformity of cob color, anther and silk color, and plant and ear height. Selection was also done for general and specific combining ability for yield in hybrid combinations across several inbreds.

From 1998 to the present, the inbred line has been observed in Bloomington, IL; Glidden, IA; Seward, NE; Washington, IA; and other locations. No phenotypic or isozymic variants have been observed.

Syngenta Seeds, Inc. NP2467 Exhibit A (Continued)

NP2467 is a uniform and stable inbred (from 2000 to 2004 during at least 6 generations of propagation).

Syngenta Seeds, Inc. NP2467 Exhibit B

Distinctness of Corn Inbred Line NP2467

The corn inbred line NP2467 (seed source 2641M60F15) us is most similar to the PVP Standard Inbred line B73 (seed source DY191083). Comparison of the two varieties was conducted in "side-by-side" trials in 2003 and 2004 at four different sites. The trial locations were Rochelle, IL; Glidden, IA Seward, NE and Bloomington, IL. The trials had two replications at each site. Plot size was 152 x 518 cm. Each plot had approximately 70 plants.

Several tables have been included in Exhibit D - Optional Additional Description of Variety, to supply the required information for all quantitative characteristics cited as distinct differences. These additional tables will contain all the statistical information including sample size, mean, LSD (at the 95% confidence level) ANOVA, the probability The method used to calculate the statistics of the combined location summary of the traits that have distinct differences use a mean value or a single data entry per replication and does not use all the individual data points per row (or subsets). The number of actual data measurements taken per quantitative traits was based upon the recommendations of the Plant Variety Protection Office. Exhibit D -Table 4 or the 2003 and 2004 Data - NP2467 vs. B73 Comparisons, contains the grand mean, number of trials with data, number of entries with data, LSD (95% confidence level), CV%, probability %, and the individual location means for all traits that are significantly different. Exhibit D - Table 5, or the 2003 and 2004 Data - NP2467 vs. B73 Comparisons – ANOVA Tables, contains additional statistical data for the traits that are significantly different. Exhibit D – Table 6 or the 2003 and 2004 Data - NP2467 vs. B73 Comparisons, Individual Replication Data, contains the data collected at all locations in 2003 and 2004 for each trait that is significantly different. Exhibit D – Table 7 or the 2003 Data - NP2467 vs. B73 Comparisons, contains the grand mean, number of trials with data, number of entries with data, LSD (95% confidence level), CV%, and probability % for the data collected at all locations in 2003 for each trait that is significantly different. Exhibit D - Table 8 or the 2004 Data - NP2467 vs. B73 Comparisons, contains the grand mean, number of trials with data, number of entries with data, LSD (95% confidence level), CV%, and probability % for the data collected at all locations in 2004 for each trait that is significantly different.

In each of these plots the B73 comparison line had a cob color at harvest of red-14.

(Munsell 10R 5/8)

In each of these plots NP2467 had a cob color at harvest of white-19. (Munsell N9.5)

NP2467 is a distinct and unique inbred line for a number of different traits but the above identified cob color easily shows the clear distinction between the two lines.

5

EXHIBIT B NP2467

	TID A LITTO			
PLOT	TRAITS	EXTERNAL	177	19
NUMBER		INBRED		BC
TOMBER	TRIAL	CODE		
			VALUE	COLOR
1011	JT610	NP2467	19	WHITE
2002	JT610	NP2467	19	WHITE
	Loc. Mean		Britis Statemen	
1007	JT610	NP2467		19
2005	JT610	NP2467		19
	Loc, Mean	a selon in it in it.		
1001	JT610	NP2467	WHITE	19
2021	JT610	NP2467	WHITE	19
	Loc, Mean			
1015	610	NP2467		19
2011	610	NP2467		19
azakar Abubahar	Loc. Mean			
1005	JT610	NP2467	19	White
2003	JT610	NP2467	19	White
	Loc. Mean			His And High I the Court of the
1005	JT610	NP2467		19
2021	JT610	NP2467		
	Loc. Mean			
1005	JT610	NP2467	WHITE	19
2029	JT610	NP2467	WHITE	19
	Loc. Mean			
1005	610	NP2467		19
2021	610	NP2467		19

EXHIBIT B B73

,	TRAITS			
PLOT	IIIII	EXTERNAL	F	219
NUMBER		INBRED	1	OBC
	TRIAL	CODE		-
			VALUE	COLOR
1002	JT610	B73	10R 7/4	8
2020	JT610	B73		
	Loc.			
	Mean	Section 19 Control 1980 (1970) and place and p		
1010	JT610	B73	10R 4/10	14
2014	JT610	B73	10R 5/8	12
	Loc.			
4040	Mean	D.7.3	2 5 MD (/ /2	
1010	JT610	B73	2.5YR 6/8	11
2006	JT610	B73	10R 5/8	14
	Loc. Mean			
1021	610	B73	10R5/8	14
2002	610	B73	10R6/8	12
	Loc.			
	Mean			
1027	JT610	B73	10	10R5/8
2005	JT610	B73	10	10R5/8
	Loc. Mean			enda en en en en en Sanda els en en en en
1027	JT610	В73	10R 5/6	14
2030	JT610	B73		
	Loc.			
1027	Mean JT610	B73	10R 5/8	14
2018	JT610	B73	10R 6/6	11
2010	Loc	D /3	1016 0/0	11
	Mean			arakas di Prilis Gult Sali Sali
1027	610	B73	10R 5/8	13
2003	610	B73	10R 5/8	13

United States Department of Agriculture, Agricultural Marketing Service Science Division, Plant Variety Protection Office National Agricultural Library Building, Room 500 Beltsville MD 20705

OBJECTIVE DESCRIPTION OF VARIETY CORN (Zea mays L.)

Name of Applicant(s) Syngenta Seeds, Inc. Address (Street & No., or R.F.D. No City, State, Zip Code and Country) Variety Seed Source 2641M60F15 Variety Name or Temporary Designat NP2467 FOR OFFICIAL USE	ion							
P.O. Box 959, Minneapolis, MN 55440 USA								
PVPO Number	1A							
Place the appropriate number that describes the varietal characters typical of this inbred variety in the spaces below. Right his thy whole numbers by adding leading zeroes if necessary. Completeness should be striven for to establish an adequate variety description. Traits designated by a '*' are considered necessary for and adequate variety description and must be completed.	h #20							
COLOR CHOICES (Use in conjunction with Munsell color code to describe all color choices: describe #25 and #26 in comment section):								
01=Light Green06=Pale Yellow11=Pink16=Pale Purple21=Buff02=Medium Green07=Yellow12=Light Red17=Purple22=Tan								
03=Dark Green 08=Yellow-Orange 13=Cherry Red 18=Colorless 23=Brown								
04=Very Dark Green 09=Salmon 14=Red 19=White 24=Bronze								
05=Green-Yellow 10=Pink-Orange 15=Red & White 20=White Capped 25=Variegated (Describe) 26=Other (Describe)								
STANDARD INBRED CHOICES(Use the most similar(in background and maturity)of these to make comparisons based on grow-out								
trial data): Yellow Dent Families: Yellow Dent (Unrelated): Sweet Corn:								
Family Members Co109, ND246, C13, Iowa5125, P39, 2132								
B14 CM105, A632, B64, B68 Oh7, T232								
B37 B37, B76, H84 W117, W153R Popcorn:								
B73 N192, A679, B73, NC268 W182BN SG1533, 4722, HP301, HP7211								
C103 Mo17, Va102, Va35, A682								
Oh43 A619, MS71, H99, Va26 White Dent: Pipecorn:								
WF9 W64A, A554, A654, Pa91 CI66, H105, Ky228 Mo15W, Mo16W, Mo24W 1. TYPE: (describe intermediate types in Comments section)								
*2 1=Sweet 2=Dent 3=Flint 4=Flour 5=Pop 6=Ornamental 7=Pipecorn								
2. REGION WHERE DEVELOPED IN THE U.S.A.: Standard Seed Source DY191083								
*5 1=Northwest 2=Northcentral 3=Northeast 4=Southeast 5=Southcentral								
6=Southwest 7=Other								
3. MATURITY (In Region Best Adaptability; show Heat Unit formula in "Comments" section):								
DAYS HEAT UNITS DAYS HEAT UNITS								
* 067 1371.0 From emergence to 50% of plants in silk 66 1344.4								
* 065 1341.8 From emergence to 50% of plants in pollen 63 1282.0	\							
003 83.5 From 10% to 90% pollen shed 002 82.1								
(*) NA NA From 50% silk to optimum edible quality NA NA								
053 924.7 From 50% silk to harvest at 25% moisture 051 1035.0								
4. PLANT:								
Standard Deviation Sample Size Standard Deviation Sample Size 212.6 cm Plant Height (to tassel tip) 16.6 16 215.3 20.4	16							
* 74.3 cm Ear Height (to base of top ear node) 9.9 16 90.0 11.9	16							
11.4 cm Length of Top Ear Internode 1.3 16 12.8 1.3	16.							
0.4 Average Number of Tillers 0.9 16 0.1 0.4	16							
* 1.2 Average Number of Ears per Stalk 0.2 16 1.4 0.3	16							
* 1.2 Average Number of Ears per Stalk 0.2 16 1.4 0.3 1								
4 Anthocyanin of Brace Roots: 1=Absent 2=Faint 3=Moderate 4=Dark Application Variety Data Page 1 Standard Inbred Data								

Δ.	onlicent M	arioty Data	Dane a	ta energia en esperante en entre en la esperante producement processa y transcripto en	Charden	Labrad Data	
5.		ariety Data	Page 2 Standard Deviation	Sample Size	Statituard	Inbred Data Standard Deviation	Sample Size
*	008.3	cm Width of Ear Node Leaf	0.7	16	009.2	0.4	16
*	078.4	cm Length of Ear Node Leaf	5.9	16	080.2	7.1	16
*	007	Number of leaves above top ear	0.6	16	006	0.4	16
	015	degrees Leaf Angle (measure from 2 nd leaf above ear at a	5.5 nthesis to stalk above lea	16 nf)	020	6.3	16
*	03	Leaf Color (Munsell code 5GY 4/4)			04	(Munsell code 5G)	(3/4)
	6	Leaf Sheath Pubescence (Rate on sca	le from1+none to 9=like	peach fuzz)	5		
	4	Marginal Waves (Rate on scale from 1	=none to 9=many)		4		
	3	Longitudinal Creases (Rate on scale fr	rom 1=none to 9=many)		4		
6.	TASSEL						
*	07	Number of Primary Lateral Branches	1.6	16	07	0.9	16
	028	Branch Angle from Central Spike	16.2	16	016	8.6	16
*	42.4	Cm Tassel Length (from top leaf collar to tassel tip)	3.0	16	39.1	2.7	16
	6	Pollen Shed (Rate on scale from 0=ma	le sterile to 9=heavy she	ed)	6		
	05	Anther Color (Munsell code 2.5GY 8/6)		26	(Munsell code) E See
	05	Glume Color (Munsell code 2.5GY 8/6))	1	26	(Munsell code)	← Comment
	2	Bar Glumes (Glume Bands): 1=Abser	nt 2=Present		2		Sel Comments Section
7a	. EAR (Un	husked Data):				· · · · · · · · · · · · · · · · · · ·	1/ Ros
*	05	Silk Color (3 days after emergence) (M	lunsell code 2.5GY 8/6)		26	(Munsell code) details
	05	Fresh Husk Color(25 days after 50% si	lking) (Munsell code 5GY	7/8)	05	(Munsell code 5G)	(6/6)
	22	Dry Husk Color (65 days after 50 % sil	king) (Munsell code 5Y 8	/6)	22	(Munsell code 5Y	8/6)
*	1	Position of Ear at Dry Husk Stage: 1=	Upright 2=Horizontal 3=	Pendent	1		
	5	Husk Tightness (Rate on scale from1=	very loose to 9=very tigh	nt)	6		
	2	Husk Extension (at harvest): 1=Short 3=Long (8-10 cm bey	(ears exposed) 2=Mediu yond ear tip) 4=Very long	ım (<8cm) g (>10 cm)	3	1	
7b *	. EAR (Hu 14.2	usked Ear Data): cm Ear Length	Standard Deviation 0.7	Sample Size 16	13.2	Standard Deviation 1.1	Sample Size
*	42.9	mm Ear Diameter at mid-point	1.4	16	43.9	2.0	. 16
	108.9	gm Ear Weight	19.7	16	98.0	23.4	16
*	16	Number of Kernel Rows	0.9	16	17	0.7	16
	2	Kernel Rows: 1=Indistinct 2=Distinct			2		
	2	Row Alignment: 1=Straight 2=Slightly	Curved 3=Spiral		1		
	6.9	cm Shank Length	1.8	16	6.8	2.2	15
	2	Ear Taper: 1=Slight 2=Average 3=Ext	reme		1		
		Application Variety Data Note: Use chart on first page to choose	e color codes for color tr	aitc		Standard Inbred D	Data

-						20034	<i>lu lu l</i>	<i>0</i>
		riety Data	Charles Deviation		Standard	Inbred Data	Carrier Clar	amendadadada
8. F		(Dried):	Standard Deviation	Sample Size		Standard Deviation	Sample Size	. ·-
	10.9	mm Kernel Length	0.5	16	11.0	0.5		15
	07.2	mm Kernel Width	0.5	16	07.4	0.8		15
	04.0	mm Kernel Thickness	0.5	16	04.2	0.5		15
	29.8	% Round Kernels (Shape Grade)	15.8	16	32.4	19.2		15
	1	Aleurone Color Pattern: 1=Homozyg	ous 2=Segregating		1			
(*)	18	Aleurone Color (Munsell code)			18	(Munsell code)	
*	07	Hard Endosperm Color (Munsell code	2.5Y 7/10)		- 06	(Munsell code 2.5Y 8/10)		
*	3	Endosperm Type: 1=Sweet (su1) 2= 4=High Amylose Starch 5=Waxy Star 8=Super Sweet (se) 9=High Oil 10=0	rch 6=High Protein 7=H		3	<i>(, -3,</i>		
	25.0	gm Weight per 100 Kernels (unsized	sample) 2.5	16	23.4	3.3		15
9. 0	ОВ		Standard Deviation	Sample Size		Standard Deviation	Sample Size	
*	26.2	mm Cob Diameter at mid-point	2.4	16	28.4	2.0	5120	15
	19	Cob Color (Munsell code N9.5)	÷		14	(Munsell code 10R	5/8)	
10.	DISEAS	E RESISTANCE (Rate from 1 (most sus	sceptible) to 9 (most res	sistant):				
	lea	ve blank if not tested: leave Race or shifts, Wilts, and Local Infection Disease	Strain Options blank if p					
В.	6	Anthracnose Leaf Blight (Colletotrichus Common Rust (Puccinia sorghi) Common Smut (Ustilago maydis) Eyespot (Kabatiella zeae) Goss's Wilt (Clavibacter michiganense Gray Leaf Spot (Cerscopora zeae-may Helminthosporium Leaf Spot (Bipolari Northern Leaf Blight (Exserohilum tur Southern Leaf Blight (Bipolaris maydis Southern Rust (Puccinia polysora) Stewart's Wilt (Erwinia stewartii) Other (Specify)	e spp. <i>Nebraskense</i>) vdis) is zeicole) Race cicum) Race		4 3	Race Race Race O		
	Stalk Ro	Corn Lethal Necrosis (MCMV and MDM Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosiac Virus (MDMV) Str Sorgham Downy Mildew of Corn (Pero Other (Specify)	ain nosclerospora soghi) n graminicola) rdis)			Strain		
Ear a	ind Kern	Gibberella Stalk Rot (Gibberella zeae) Other (Specify) el Rots Aspergillus Ear and Kernel Rot (Aspergipolis Ear Rot (Stenocarpella maydifusarium Ear and Kernel Rot (Fusarium Gibberella Ear Rot (Gibberella zeae)	gillus flavus) is)					
Annli		Other (Specify) ariety Data		·	Ctondond	Inhrad Data		
Note	Use ch	arrety bata lart on first page to choose color codes	s for color traits		Stanual U	Inbred Data		
				· · · · · · · · · · · · · · · · · · ·				

Butler, D.R. 1954. A System for the Classification of corn Inbred Lines. PhD thesis. Ohio State University.

Emerson, R. A., G.W. Beadle, and A.C. Fraser. 1935. A Summary of Linkage Studies in Maize. Cornell A.E.S., Mem. 180

Farr, D.F., G.F. Bills, G.P. Chamuris, A.Y. Rossman. 1989. Fungi on Plant and Plant Products in the United States. The American Phytopathological Society, St. Paul, MN.

Inglett, G.E. (Ed) 1970. Corn: Culture, Processing, Products. Avi Publishing Company, Westport, CT.

Jugenheimer, R.W. 1976. Corn: Improvement. Seed Production, and Uses. John Wiley & Sons, New York.

McGee, D.C. 1988. Maize Diseases. AOS Press, St. Paul, MN. 150 pp.

Munsell Color Chart for Plant Tissues. Mcbeth. P.O. Box 230. Newburgh, N.Y. 12551-0230

The Mutants of Maize. 1968. Crop Science Society of America. Madison, WI.

Shurtleff, M.C. 1980. Compendium of Corn Diseases. APS Press, St. Paul, MN. 105 pp.

Sprague, G.F., and J.W. Dudley (Editors). 1988 Corn and Corn Improvement, Third Edition. Agronomy Monograph 18. ASA, SCCA, SSSA,

Stringfield, G.H, Maize Inbred Lines of Ohio. Ohio A.E.S., Bul. 831. 1959

U.S. Department of Agriculture. 1936, 1937. Yearbook.

COMMENTS (eg. state how heat units were calculated, standard inbred seed source, and /or where data was collected. Continue in Exhibit

Heat Units per day were calculated using the standard formula: HU={MaxTemp (86) + Min Temp (50)]/2-50. 1)

Data for this exhibit was collected at London, Ontario, Canada, Stanton, MN and Janesville, WI.

Large standard deviations are probably due to environmental factors at each individual location where the variety was observed. Since the varieties reported in this exhibit are inbreds, the response to the environment is probably more pronounced than a hybrid or a combination of these inbred lines. Any stress at specific times during the growing season could influence results.

The Anther color of B73 is green yellow (2.5GY 8/6 or 05) with purple (17).

- The glume color of B73 is green yellow (2.5GY 8/6 or 05) with purple (17). Silk Color of B73 is green/yellow (2.5GY 8/6 or 05) but shades pale purple (2.5R 6/4 or 16) with age.

The Insect data reported was collected at Bloomington, IL in 2004

8) The Disease data reported was collected at Bloomington, IL in 2004


Point 5(a, (b) and (c)-There is a concern that there are large Standard deviations for certain NP2467 and B73 data. As noted by the Examiner the Trial conditions of the locations may have been the cause of this. The environmental data for London Ontario Canada and for the Stanton Minnesota trials in the two years is being submitted. The environmental information for Janesville is not presently available.

The London trial location may be causing the standard deviations for these plants to be a bit wider than normal for a couple of reasons. The planting dates for the London trials was substantially later than the dates for the Stanton planting. London is in an area that plants 83-90 day corn and Stanton and Janesville are locations that plant 94-102 day corn. In addition to the relative maturity difference between the locations, the delayed plantings due to the weather conditions in London provide even less time

for the plants to mature.


London normally plants in the May 5- May 10 range. But in both years the actual planting dates were substantially later than this date. The environment in 2003 was hotter than normal. To the extent that London trial resorted to nursery irrigation to attempt to avoid moisture stress at flowering time. In contrast, London had a very wet spring in 2004 which even further delayed the already late dates for planting. If the unusual environmental conditions are removed from the Exhibit C data it is believed that the standard deviations for the standard line B73 and NP2467 would be more in line with expected numbers. If the Examiner would like a new Exhibit C with the outlier data from out of area trials removed and the standard deviations for the Exhibit recalculated we would request that the Examiner contact us so that we can supply the substitute Exhibit C. Thanks for your review of this information. Please contact us if you need further information for your review of the application.

Additional Description of the Corn Inbred Line NP2467 Table 1




Typical B73 Silk and Second Leaf Color

Additional Description of the Corn Inbred Line NP2467 Table 1 (continued)




Typical NP2467 Silk and Second Leaf Color

Additional Description of the Corn Inbred Line NP2467 Table 2




Typical B73 Tassel

Additional Description of the Corn Inbred Line NP2467 Table 2 (continued)



Typical NP2467 Tassel

Additional Description of the Corn Inbred Line NP2467 Table 3



Anther and Glume Color of B73



Anther and Glume Color of NP2467

2003 and 2004 Data - NP2467 vs. B73 Comparisons Syngenta Seeds, Inc. Exhibit D - Table 4

Combined Summary:

NP2467

LSD General (5%) EE CV (Effective) % Probability % Entries w/data Mean General Mean Control Trials w/data

Individual Location Data - NP2467: 2003 Bloomington, IL 2004 Bloomington, IL 2004 Glidden, IA 2004 Seward, NE 2003 Seward, NE 2003 Rochelle, IL 2004 Rochelle, IL 2003 Glidden, IA

Mean

ndividual Location Data - B73: 2004 Bloomington, IL 2003 Bloomington, IL 2003 Seward, NE 2004 Seward, NE 2003 Rochelle, IL 2004 Rochelle, IL 2003 Glidden, IA 2004 Glidden, IA

Ear	Weight	(am)
Ear	Height	(cm)
Cop	Diameter	(mm)
Angle	2nd	<u>Leaf</u>

			_				_	
109	86	103		8	2	6	7	3
74	06	82		8	7	7	7	0
	28	27		8	2	1	3	0
15	20	18		8	2	4	21	3

00			
07		54	
20		74	
10		69	66
6	28	82	
20	24	85	
8	25	72	
18		90	
18		80	
15	26	74	

	57	66	66	8	101	141	108	97	86
	63	92	91	100	101	88	68	26	06
	28	26	30	31	28	26	29	31	28
	20	20	20	10	20	15	30	28	20
-									

Syngenta Seeds, Inc. 2003 and 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 4

Combined Summary:

NP2467 B73 Mean General
Mean Control
Trials w/data
Entries w/data
LSD General (5%) EE
CV (Effective) %
Probability %

Individual Location Data - NP2467:

2003 Rochelle, IL 2003 Glidden, IA 2003 Seward, NE 2003 Bloomington, IL 2004 Rochelle, IL 2004 Glidden, IA 2004 Seward, NE

2004 Bloomington, IL Mean Individual Location Data - B73:

2003 Rochelle, IL 2003 Glidden, IA

2003 Seward, NE 2003 Bloomington, IL

2004 Rochelle, IL

2004 Glidden, IA 2004 Seward, NE

2004 Bloomington, IL Mean

Lengt Tass								
Heat Units to 50% Pollen Shed	1342	1282	1312	80	2	27	2	-
Heat Units to 90% <u>Pollen Shed</u>	1389	1316	1353	8	2	46	3	
Heat Units to 10% Pollen Shed	1321	1249	1285	7	2	26	2	•

42	1342	1389	1321
42	1244	1276	
37	1257	1296	1227
42	1327	1351	1306
43	1193	1209	1169
41	1642	1678	1616
46	1370	1450	1334
46	1321	1345	1294
41	1385	1507	1304

35	43	43	39	39	39	40	36	39
1266	1292	1302	1589	1124	1249	1243	1196	1282
1318	1317	1346	1630	1181	1263	1267	1209	1316
1213	1267	1232	1549	1069	1233	1185		1249

2003 and 2004 Data - NP2467 vs. B73 Comparisons Syngenta Seeds, Inc. Exhibit D - Table 4

Combined Summary:

Stay Length of Internode (cm)

Rating Green

5	2	က	9	2	2	46	0
1	13	12	8	2		6	2

						_		
ဍ		5	2	4		က	7	5
6	11	12	11	12	12	13	12	17

Individual Location Data - NP2467:

2003 Rochelle, IL

2003 Bloomington, IL

2004 Rochelle, IL 2004 Glidden, IA

2003 Seward, NE

2003 Glidden, IA

LSD General (5%) EE

Entries w/data

Mean General Mean Control Trials w/data

NP2467

CV (Effective) %

Probability %

2		1	3	-		7	2	2
11	14	14	13	14	14	11	12	13

Individual Location Data - B73:

2003 Rochelle, IL

2003 Bloomington, IL

2004 Rochelle, IL

2003 Seward, NE

2003 Glidden, IA

2004 Bloomington, IL

Mean

2004 Seward, NE

2004 Glidden, 1A

2004 Bloomington, IL

Mean

2004 Seward, NE

Syngenta Seeds, Inc. 2003 and 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 5

TRAIT	SOURCE	DF	SS1	SS2	MS	L ,	P_PCT
Angle 2nd <u>Leaf</u>	Treatment Location Treatment*Location Pooled Error Total	1 7 7 185 15	105.42 386.49 96.66 588.57	105.42	105.42 55.21 13.81 15.81	7.63	2.7977
Cob Diameter (mm)	Treatment Location Treatment*Location Pooled Error Total	1 7 7 179 15	18.93 59.78 6.35 85.06	18.93	18.93 8.54 0.91	9.41	0.2581
Ear Height <u>(cm)</u>	Treatment Location Treatment*Location Pooled Error Total	1 7 7 190 15	984.93 1610.89 69.89 2665.71	984.93	984.93 230.13 9.98 26.35	98.65 23.05	0.0022
Ear Weight <u>(gm)</u>	Treatment Location Treatment*Location Pooled Error Total	1 7 7 7 179 115	456.87 6146.47 399.98 7003.32	456.87	456.87 878.07 57.14 79.62	15.37	2.5492 0.093
Heat Units to 10% <u>Pollen Shed</u>	Treatment Location Treatment*Location Pooled Error Total	6 6 158 13	18126.59 246920.06 2459.49 267506.13	18126.59	18126.59 41153.34 409.92 1154.82	44.22 100.39	0.0559

Syngenta Seeds, Inc. 2003 and 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 5

TRAIT	SOURCE	DF	SS1	SS2	MS	<u>ш</u>	P_PCT
Heat Units to 90% Pollen Shed	Treatment Location Treatment*Location Pooled Error Total	1 7 7 186 15	21133.52 281391.79 10702.43 313227.74	21133.52	21133.52 40198.83 1528.92 1047.07	13.82 26.29	0.7477
Heat Units to 50% Pollen Shed	Treatment Location Treatment*Location Pooled Error Total	1 7 7 186 15	14278.43 258520.85 3609.91 276409.19	14278.43	14278.43 36931.55 515.7 1023.91	27.69	0.1171
Length of Tassel (<u>cm)</u>	Treatment Location Treatment*Location Pooled Error Total	1 7 7 161 15	42.03 90.58 28.79 161.4	42.03	42.03 12.94 4.11 6.36	3.15	1.5127 7.6717
Length of Internode (<u>cm)</u>	Treatment Location Treatment*Location Pooled Error Total	1 7 7 190 15	7.78 14.33 9.2 31.3	7.78	7.78 2.05 1.31 1.25	5.92 1.56	4.5229 28.652
Stay Green <u>Rating</u>	Treatment Location Treatment*Location Pooled Error Total	- 2 4 - 2 5 1 - 1	28.79 6.63 14.01 49.44	28.79	28.79 1.33 2.8 0.49	10.27 0.47	2.3866 78.4358

Syngenta Seeds, Inc. 2003 and 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 6 - Individual Replication Data

4																										
·	28	25	26	27	3 6	24	26	27	26	29	27	28	25	24	24	25	25	25	27	23	25	31	30	31	26	2.4
L	28			25	23	2		26		30	30		25	25		26	24		26	23		32	32			
Cob Diameter (mm)	27	i		24	23		24	27		8	25		23	24		25	24		27	23		30	30			
Dis.	28			24	23		24	27		8	27		24	24		78	24		28	24		8	30			
	28	25		24	23		27	28		29	26		26	24		25	26		26	22		33	30			
	28		<u> </u>	24		<u> </u>	28			28	22		25	24		23	26		27	23		32	30			
Angle 2nd <u>Leaf</u>	20	20	20	15	25	20	10	10	10	7	10	6	20	20	20	5	10	8	16	20	18	20	15	18	15	5.5
Year <u>Date</u>	2003	2003		2003	2003		2003	2003		2003	2003		2004	2004		2004	2004		2004	2004		2004	2004	-		
Replication	Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2			
<u>Loc.</u>	Rochelle, IL	Rochelle, IL	Location Mean	Glidden, IA	Glidden, IA	Location Mean	Seward, NE	Seward, NE	Location Mean	Bloomington, IL	Bloomington, IL	Location Mean	Rochelle, IL	Rochelle, IL	Location Mean	Glidden, IA	Glidden, IA	Location Mean	Seward, NE	Seward, NE	Location Mean	Bloomington, IL	Bloomington, IL	Location Mean		
Inbred <u>Name</u>	NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		Mean	Std. Dev.

Syngenta Seeds, Inc. 2003 and 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 6 - Individual Replication Data

	93	73	83	3 8	95	8 8	66	66	66	100	66	66	116	122	119	140	157	149	107	107	107	104	130	117	109	19.7
	94	1		107	5			190		86	118		112	136		164	151		104	100		110	149			
Ear Weight (gm)	55			Ę	66		54	66		66	137		104	113		112	140		111	103		116	132			
- × 3	84			Ę	56	3	66	104		66	51		120	112		125	177		105	104	-	120	139			
	117	80		105	93		119	97	!	103	97		127	128		159	130		110	120		84	130			
	119	65		100	96		125	103		101	97		119	120		142	185		107	110		6	86			
	49	28	54	12	72	74	62	75	69	84	79	82	91	78	85	20	75	72	84	92	80	79	8	80	74	6.6
	50	70		74	99		62	89		83	78		06	80		92	89		81	73		70	70			
Ear Height <u>(cm)</u>	45	09		88	74	-	20	70		101	80		90	80		70	72		84	77		75	100			
H 위	50	55		2	70		65	98		80	80		95	65		70	92		76	75		85	80			
	20	20		76	70	 	09	83		75	83		06	06		72	99		85	78		80	75			
	20	55		9/	78		75	70		80	75		90	75		09	74		92	76		85	75			
Year Date	2003	2003		2003	2003		2003	2003		2003	2003		2004	2004		2004	2004		2004	2004		2004	2004			
Replication	Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2			
<u>Loc.</u>	Rochelle, IL	Rochelle, IL	Location Mean	Glidden, IA	Glidden, IA	Location Mean	Seward, NE	Seward, NE	Location Mean	Bloomington, IL	Bloomington, IL	Location Mean	Rochelle, IL	Rochelle, IL	Location Mean	Glidden, IA	Glidden, IA	Location Mean	Seward, NE	Seward, NE	Location Mean	Bloomington, IL	Bloomington, IL	Location Mean		The state of the s
Inbred <u>Name</u>	NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		Mean	Std. Dev.

Syngenta Seeds, Inc. 2003 and 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 6 - Individual Replication Data

Loc. Rochelle, IL		ī			
Rochelle, IL	Replication	Date	Pollen Shed	Pollen Shed	to 50% Pollen Shed
	Rep. 1	2003	1362	1570	1408
Rochelle, IL	Rep. 2	2003	1245	1435	1361
Location Mean			1304	1507	1385
Glidden, IA	Rep. 1	2003	1294	1345	1204
Glidden, IA	Rep. 2	2003			1 701
Location Mean			1294	1345	1201
Seward, NE	Rep. 1	2003	1334	1450	1358
Seward, NE	Rep. 2	2003	1334	1450	1381
Location Mean			1334	1450	1370
Bloomington, IL	Rep. 1	2003	1603	1678	1630
Bloomington, IL	Rep. 2	2003	1630	1678	1654
Location Mean			1616	1678	1642
Rochelle, IL	Rep. 1	2004	1169	1209	1193
Rochelle, IL	Rep. 2	2004	1169	1209	1193
Location Mean			1169	1209	1193
Glidden, IA	Rep. 1	2004	1306	1351	1327
Glidden, IA	Rep. 2	2004	1306	1351	1327
Location Mean			1306	1351	1327
Seward, NE	Rep. 1	2004	1233	1301	1257
Seward, NE	Rep. 2	2004	1221	1291	1257
Location Mean			1227	1296	1257
Bloomington, IL	Rep. 1	2004		1288	1278
Bloomington, IL.	Rep. 2	2004		1264	1209
Location Mean				1276	1244
			1321	1389	1342
			141.8	150.4	137.5

Syngenta Seeds, Inc. 2003 and 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 6 - Individual Replication Data

Name	<u>Loc.</u>	Replication	Year <u>Date</u>			Tassel (cm)	sel (i)			of Internode (cm)
NP2467	Rochelle, IL	Rep. 1	2003	44	52	42	42	42	44	10
NP2467	Rochelle, IL	Rep. 2	2003	40	37	35	39	38	38	0 &
	Location Mean								41	0
NP2467	Glidden, IA	Rep. 1	2003	56	28	36	46	48	49	9
NP2467	Glidden, IA	Rep. 2	2003	48	42	42	42	46	44	12
	Location Mean								46	7 7
NP2467	Seward, NE	Rep. 1	2003	47	26	36	4	20	46	- 2
NP2467	Seward, NE	Rep. 2	2003	44	50	46	45	48	47	1
	Location Mean								46	12
NP2467	Bloomington, IL	Rep. 1	2003	43	46	36	14	38	4	
NP2467	Bloomington, IL	Rep. 2	2003	39	4	44	46	40	42	10
	Location Mean				·				4	7
NP2467	Rochelle, IL	Rep. 1	2004	35	41	45	42	47	42	-
NP2467	Rochelle, IL	Rep. 2	2004	41	42	47	46	40	43	13
	Location Mean								43	12
NP2467	Glidden, IA	Rep. 1	2004	44	4	4	20	38	42	12
NP2467	Glidden, IA	Rep. 2	2004							12
	Location Mean								42	12
NP2467	Seward, NE	Rep. 1	2004	39	41	35	33	38	37	12
NP2467	Seward, NE	Rep. 2	2004	37	39	33	37	36	37	14
	Location Mean								37	13
NP2467	Bloomington, IL	Rep. 1	2004	44	46	45	41	40	43	12
NP2467	Bloomington, IL	Rep. 2	2004	44	4	42	36	39	41	12
	Location Mean								42	12
Mean								-	42	
Std. Dev.									27	13

Syngenta Seeds, Inc. 2003 and 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 6 - Individual Replication Data

	r	1-2-	T		1	,	1		T	T = .			T	····	····	,		····	T ==	T	l				1	
Stay Green Rating	ပ	4	5				3	7	5	2	2	2	0	4	4				8	8	3	7	7	7	5	1.9
Year <u>Date</u>	2003	2003		2003	2003		2003	2003		2003	2003		2004	2004		2004	2004		2004	2004		2004	2004			
Replication	Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2			
<u>Loc.</u>	Rochelle, IL	Rochelle, 1L	Location Mean	Glidden, IA	Glidden, IA	Location Mean	Seward, NE	Seward, NE	Location Mean	Bloomington, IL	Bloomington, IL	Location Mean	Rochelle, IL	Rochelle, IL	Location Mean	Glidden, IA	Glidden, IA	Location Mean	Seward, NE	Seward, NE	Location Mean	Bloomington, IL	Bloomington, IL	Location Mean		
Inbred Name	NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		NP2467	NP2467		Mean	Std. Dev.

Syngenta Seeds, Inc. 2003 and 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 6 - Individual Replication Data

	28		28			26		30	99		31	31	29		28	26		26		29	29	31	31	31	28	0.0
ter (25			29			30	30		28	27		27	27		30			32	30			
Cob Diameter (mm)	3 28			25	25		30	28		30	32		28	28		25	25		28	29		30	30			
<u></u>	3 28			3 26	3 25		28	28		31	32		28	, 27		27	26		29	30		30	32			
	3 28	_		3 26	3 26		31	32		28	32		30	27		27	25		29	28		32	30			
	28		_) 26	26		30	31	_	30	30		29	28		25	25		29	28	_	30	32			
Angle 2nd <u>Leaf</u>	20	20	20	20	20	20	10	30	20	5	15	10	20	20	20	15	15	15	30	30	30	25	30	28	20	6.3
Year <u>DATE</u>	2003	2003		2003	2003		2003	2003		2003	2003		2004	2004		2004	2004		2004	2004		2004	2004			
Replication	Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2			
<u>Loc.</u>	Rochelle, IL	Rochelle, IL	Location Mean	Glidden, IA	Glidden, IA	Location Mean	Seward, NE	Seward, NE	Location Mean	Bloomington, IL	Bloomington, IL	Location Mean	Rochelle, IL	Rochelle, IL	Location Mean	Glidden, IA	Glidden, IA	Location Mean	Seward, NE	Seward, NE	Location Mean	Bloomington, IL	Bloomington, IL	Location Mean		
Inbred <u>Name</u>	B73	B73		B73	B73		B73	B73		B73	B73		B73	B73		B73	B73		B73	B73		B73	B73		Mean	Std. Dev.

Syngenta Seeds, Inc. 2003 and 2004 Data - NP2467 vs. B73 Comparisons

Exhibit D - Table 6 - Individual Replication Data

Syngenta Seeds, Inc. 2003 and 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 6 - Individual Replication Data

	Inbred	<u>Loc.</u>	Replication	Year DATE	Heat Units to 10% Pollen Shed	Heat Units to 90% Pollen Shed	Heat Units to 50% Pollen Shed
	B73	Rochelle II	Ren 1	2003	1999	1326	000
	B73	Rochelle, IL	Rep. 2	2003	1204	1340	1200
<u> </u>		Location Mean			1213	1318	1266
	B73	Glidden, IA	Rep. 1	2003	1241	1292	1265
	B73	Glidden, IA	Rep. 2	2003	1292	1342	1318
		Location Mean			1267	1317	1292
	B73	Seward, NE	Rep. 1	2003	1245	1358	1314
	B73	Seward, NE	Rep. 2	2003	1219	1334	1290
		Location Mean			1232	1346	1302
	B73	Bloomington, IL	Rep. 1	2003	1549	1630	1603
	B73	Bloomington, IL	Rep. 2	2003	1549	1630	1575
		Location Mean			1549	1630	1589
	B73	Rochelle, IL	Rep. 1	2004	1079	1193	1145
	B73	Rochelle, IL	Rep. 2	2004	1058	1169	1103
l		Location Mean			1069	1181	1124
	B73	Glidden, IA	Rep. 1	2004	1224	1256	1241
	B73	Glidden, IA	Rep. 2	2004	1241	1270	1256
		Location Mean			1233	1263	1249
	B73	Seward, NE	Rep. 1	2004	1198	1270	1251
	B73	Seward, NE	Rep. 2	2004	1172	1264	1235
		Location Mean			. 1185	1267	1243
	B73	Bloomington, IL	Rep. 1	2004		1209	1196
	B73	Bloomington, IL	Rep. 2	2004		1209	1196
[Location Mean				1209	1196
	Mean			(1250	1316	1282
	Std. Dev.	prophasis () and ()			146.5	138.4	136.2

Syngenta Seeds, Inc. 2003 and 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 6 - Individual Replication Data

Rep. 1 2003 39 Rep. 2 2003 36 Rep. 1 2003 46 Rep. 2 2003 34 Rep. 1 2003 39 Rep. 2 2003 47 Rep. 1 2003 41 Rep. 2 2003 41 Rep. 1 2003 41 Rep. 2 2004 36 Rep. 2 2004 43 Rep. 2 2004 43
1 2003 2 2003 1 2003 2 2003 1 2003 2 2003 1 2004 2 2004
2 2003 2 2003 1 2003 2 2003 1 2003 2 2003 2 2004 2 2004
1 2003 1 2003 1 2003 2 2003 2 2004 1 2004 2 2004
1 2003 1 2003 2 2003 1 2003 2 2003 2 2004 2 2004
2 2003 1 2003 2 2003 1 2003 2 2004 2 2004
2 2003 1 2003 2 2003 2 2004 2 2004
1 2003 2 2003 1 2003 2 2004 1 2004 2 2004
2 2003 1 2003 2 2003 1 2004 2 2004
1 2003 2 2003 1 2004 2 2004
2 2003 1 2004 2 2004
2 2003 1 2004 2 2004
2 2004
2 2004
2 2004
Rep. 1 2004 44
Rep. 2 2004
Rep. 1 2004 40
Rep. 2 2004 41
Rep. 1 2004 36
Rep. 2 2004 36
The state of the s

Syngenta Seeds, Inc. 2003 and 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 6 - Individual Replication Data

Stay Green	ating	2	2	2				7	_	-	2	4	3	1	-	1				1	1	1	1	2	2	2	0.8
	DATE	2003	2003		2003	2003		2003	2003		2003	2003		2004	2004		2004	2004		2004	2004		2004	2004			
	Replication	Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		Rep. 1	Rep. 2		<u></u>	
	<u>Loc.</u>	Rochelle, IL	Rochelle, IL	Location Mean	Glidden, IA	Glidden, IA	Location Mean	Seward, NE	Seward, NE	Location Mean	Bloomington, IL	Bloomington, IL	Location Mean	Rochelle, IL	Rochelle, IL	Location Mean	Glidden, IA	Glidden, IA	Location Mean	Seward, NE	Seward, NE	Location Mean	Bloomington, IL	Bloomington, IL.	Location Mean		
Inbred	Name	B73	B73		B73	B73		B73	B73		B73	B73		B73	B73		B73	B73	·	B73	B73		B73	B73		Mean	Std. Dev.

Syngenta Seeds, Inc. 2003 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 7

Combined Summary:

NP2467 B73 Mean General
Mean Control
Trials w/data
Entries w/data
LSD General (5%) EE
CV (Effective) %
Probability %

Ear Weight (<u>gm)</u>	95	85	06	4	2		10	20
Ear Height (cm)	69	86	78	4	2	8	5	1
Cob Diameter (mm)	26	28	27	4	2		2	1
Angle 2nd <u>Leaf</u>	15	18	16	4	2		21	32

Syngenta Seeds, Inc. 2003 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 7

Combined Summary:

NP2467
B73
Mean General
Mean Control
Trials w/data
Entries w/data
LSD General (5%) EE
CV (Effective) %
Probability %

Length of Tassel (cm)	44	40	42	4	2	2	က	2
Heat Units to 50% Pollen Shed	1429	1362	1396	4	2	09	2	4
Heat Units to 90% <u>Pollen Shed</u>	1495	1402	1449	4	2		4	8
Heat Units to 10% Pollen Shed	1387	1315	1351	4	2	52	2	2

Syngenta Seeds, Inc. 2003 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 7

Combined Summary:

Stay Green Rating

Length of Internode (cm)

NP2467 B73

Mean General

LSD General (5%) EE CV (Effective) % Probability % Entries w/data Mean Control Trials w/data

4	7	3		2			16
11	13	12	4	2	0	1	0

Syngenta Seeds, Inc. 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 8

Combined Summary:

NP2467 B73 Mean General
Mean Control
Trials w/data
Entries w/data
LSD General (5%) EE
CV (Effective) %
Probability %

က	N	Τ	7	Г	4	2	Г	9	N
123			117						12
62	94		98		7	2	9	E	0
26	28		27		4	2		5	13
16	23		19		4	2		19	7

Ear Weight (<u>gm)</u>

Ear Height (cm)

Cob Diameter <u>(mm)</u>

Angle 2nd <u>Leaf</u>

Syngenta Seeds, Inc. 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 8

Combined Summary:

NP2467 B73 Mean General
Mean Control
Trials w/data
Entries w/data
LSD General (5%) EE
CV (Effective) %
Probability %

neat Units to 10% Pollen Shed	Heat Units to 90% Pollen Shed	Heat Units to 50% Pollen Shed	Length of Tassel (cm)
1234	1283	1255	41
1162	1230	1203	38
1198	1257	1229	40
			,
3	4	4	4
2	2	2	2
	47	46	
2	2	2	9
5	4	4	16

Syngenta Seeds, Inc. 2004 Data - NP2467 vs. B73 Comparisons Exhibit D - Table 8

Combined Summary:

Stay Green Rating

Length of Internode (cm)

NP2467 B73

Mean General Mean Control

LSD General (5%) EE Entries w/data Trials w/data

CV (Effective) % Probability %

٥ ا	13 1	12 3		2 2		9 29	
						3	

REPRODUCE LOCALLY. Include form number and edition date on al	reproductions.	FORM APPROVED - OMB No. 0581-0055
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to del certificate is to be issued (7 U.S.C. 2 confidential until the certificate is issued.	421). The information is held
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME
Syngenta Seeds, Inc.		NP2467
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (Include area code)	6. FAX (Include area code)
P.O. Box 959 Minneapolis, MN 55440	(763) 593-7333	(763) 593-7828
· · · · · · · · · · · · · · · · · · ·	7. PVPO NUMBER	
·		200500100
8. Does the applicant own all rights to the variety? Mark an "X" in the	e appropriate block. If no, please expla	ain. YES NO
9. Is the applicant (individual or company) a U.S. national or a U.S. b	, , , , , ,	
10. Is the applicant the original owner?	NO If no, please answer <u>one</u>	of the following:
a. If the original rights to variety were owned by individual(s), is ((are) the original owner(s) a U.S. Nation NO If no, give name of coun	
b. If the original rights to variety were owned by a company(ies)	, is (are) the original owner(s) a U.S. ba	
•		
11. Additional explanation on ownership (Trace ownership from origin	nal breeder to current owner. Use the r	everse for extra space if needed):
(See attached Exhibit E Addendum)		
PLEASE NOTE:		_

Plant variety protection can only be afforded to the owners (not licensees) who meet the following criteria:

- 1. If the rights to the variety are owned by the original breeder, that person must be a U.S. national, national of a UPOV member country, or national of a country which affords similar protection to nationals of the U.S. for the same genus and species.
- If the rights to the variety are owned by the company which employed the original breeder(s), the company must be U.S. based, owned by nationals of a UPOV member country, or owned by nationals of a country which affords similar protection to nationals of the U.S. for the same genus and species.
- 3. If the applicant is an owner who is not the original owner, both the original owner and the applicant must meet one of the above criteria.

The original breeder/owner may be the individual or company who directed the final breeding. See Section 41(a)(2) of the Plant Variety Protection Act for definitions.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is estimated to average 0.1 hour per response, including the time for reviewing the instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require afternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.

Syngenta Seeds, Inc. NP2467 Exhibit E, Block #11

Additional Explanation on Ownership

Syngenta breeder, Albert R. Hornbrook, developed inbred NP2467, working primarily from Boynton Beach, Florida and Bloomington, Illinois (see Exhibit A for detail). Hornbrook was a corn breeder for Ciba Geigy Company at the beginning of the line's development, was employed by Novartis Seeds, Inc. (formed by the merger of Ciba Seeds and Sandoz Seeds) during development of the inbred, and was employed by Syngenta Seeds, Inc. at the completion of the line's development in 2000. This line is exclusively owned by Syngenta Seeds, Inc. (formed by the merger of the agribusinesses of Novartis and Zeneca in 1999).